

## Original Research Article

# Standing Tree Biomass and Carbon Stock of Natural Forests in Nainital District of Kumaun Himalaya

\*Bijendra Lal, L.S. Lodhiyal and Dhan Kumar Bhatt<sup>1</sup>

College of Forestry (VCSG, UUFH) Ranichauri, Tehri Garhwal-249199

<sup>1</sup>Department of Forestry and Environmental Science, DSB Campus Kumaun University, Nainital-263002, Uttarakhand

\*Corresponding author: bijendraforestry85@gmail.com

Received: June 29, 2025; revised: August 9, 2025; accepted: August 14, 2025

<https://doi.org/10.5281/zenodo.16949675>

Published Online: September 2, 2025

**Abstract:** Present study deals with the standing tree biomass and carbon stock of natural forests in Nainital, district of Kumaun Himalaya. Uttarakhand the 27<sup>th</sup> state of the India is located between 28°43'-31°27' N latitude and 77°34'-81°02' E longitudes. Forests in the region of Uttarakhand are mainly composed of three dominated tree species such as Sal (*Shorea robusta* Gaertn.) in lower elevation (400-1200m), pine (*Pinus roxburghii* Sarg. in mid elevation (1200-1800m) and Oak (*Quercus* spp.) in higher elevation (1400-2700m), which indicates that the forest tree composition changes with the change in altitude and climate in the region. Vegetation analysis was carried out by using quadrat method in each selected forest sites. The tree species were analyzed by using quadrat size 10m x 10m in each study forest site. In each tree species were measured at 1.37m. The ecological parameters like tree density, abundance, basal area and IVI of trees estimated in each study forest site. The ecological parameters like tree density, abundance, basal area and IVI of trees estimated in each forest site as followed by Misra (1968). The tree density range from 424-796 ind.ha<sup>-1</sup>. The total basal areas of tree range between 40.41-49.42 m<sup>2</sup> ha<sup>-1</sup>. The estimation of biomass was carried out using the allometric equation for each tree component. The biomass and carbon stock of the trees ranged from 135.43-513.20 t ha<sup>-1</sup>yr<sup>-1</sup> and 64.33-243.77 t ha<sup>-1</sup>yr<sup>-1</sup>. Based on the findings it can be concluded that vegetation structure, biomass and carbon stock of natural forest need the certain management inputs for their better conservation and development.

**Keywords:** Biomass, basal area, carbon stock, Kumaun Himalaya, tree density.

## Introduction

The Indian Himalaya is a vast mountain system with a geographical area of about 591000sq.km responding 18% area of the country and is an important part of the global system. The Himalayan region is one of the most dynamic and complex mountain ranges in the world due to tectonic activity and is very vulnerable due to global climate changes. Global change whether generated by climate, land use change, biological invasion, global economic forces, will certainly affect the

relationships that are present in the land and economics of the Himalayan region. Uncertainties and potential impacts of the rate and magnitude of climate change prevail, but there is no question that it is gradually and powerfully changing the ecological and socio economic landscape of the region. The fact is that India is one of the 17<sup>th</sup> mega biodiversity regions of the world are mainly due to the Himalaya region. The Indian Himalayan region is also called the 'water tower of the earth'.

Roughly 10-20% of the area is covered by glaciers, while 30-40% remains under seasonal snow cover (Bahadur, 2004). Despite the vast water resources movements such as retreating effects of glaciers, streams and rivers occurs regularly in the region. The Himalaya is a provider of variety of natural resources to the great Gangetic plains the heartland of the country including the life giving water. Researchers believe that these resources have largely been endangered by human activities due to over exploitation of forest products, uncontrolled grazing, encroachment and deforestation on the forest area. (Champion and Seth, 1968; Singh and Singh, 1991; Eckholm, 1975).

Himalayan forest vegetation ranges from tropical dry deciduous forest in the foothills to alpine meadow above timberline (Singh and Singh, 1992). Composition of the forest is diverse and varies from place to place because of varying topography such as plains, foothills and upper mountains (Singh, 2006). Economically and environmentally, the natural resources are the main source for people in this region (Ram *et al.*, 2004). In addition, environmental problems are particularly noticeable in this region as a form of degradation and depletion of the forest resources (Sati, 2005).

The people of Indian Himalayan region resembling other mountain ecosystems, are greatly reliant for livelihood on their immediate natural resources and production from most important sectors i.e. agriculture, forestry and livestock, etc. The dependency of the ever increasing population on limited resources is getting higher. Lack of modern technology to reduce mountain specificities and enhanced production to meet the burden are exhausting the resources in conjunction with marginality of farmers in the end advancing poverty (Samant *et al.*, 2003). Despite its rich biological resources the region is underdeveloped. It won't be an understatement to call it a third world of a developing nation. Trends of environmental well being indicate that existing interventions are unsustainable. Economic indicators also do not show the desirable effects on monetary upliftment, moreover, the natural delicateness of the mountains as well as the vulnerability of the Himalaya to human induced environmental

impacts make people live in the gloom of uncertainties of natural calamities.

Biomass is defined as the total amount of AGB in trees expressed as oven-dry tons per unit area that reduces the concentration from atmospheric concentration of CO<sub>2</sub>. Biomass is an ecological material derived from living or recently living organism. As an energy source biomass can either be directly used via combustion to produce heat, or indirectly after converting it to various forms of bio-fuel. Biomass is the mass of originally bound carbon (C) that is present in forest area. Apart from bacteria, the total live biomass on earth is about 560 billion tones C and the total annual primary production of biomass is just over 100 billion tones t/yr in the study forest area. However, the total live biomass of bacteria may exceed that of plants and animal.

Anderson (1970) opined that estimation of biomass is essential for determining the status and flux of biological materials in ecosystem and for understanding the dynamics of the ecosystem. The quantity of the tree biomass per unit area of land constitutes the primary inventory data needed to understand the flow of materials and water through forest ecosystem (Swank and Schedules 1974). The primary objectives of this study is to determine different vegetation parameters available in the sites and to calculate biomass and carbon stock in each selected forests.

## Materials and methods

### Site description:

The present study was conducted in three selected dominant forests viz., Oak (*Quercus leucotrichophora*, A. Camus) dominated forest; Pine (*Pinus roxburghii*, Sarg.) dominated forest and Sal (*Shorea robusta*, Gaertn. F.) dominated forest located in Kailakhan (29° 24' N; 79° 28' E), Sanitorium (29° 23.634' N; 79° 30.466' E) and the Ranibagh (29° 17.803' N; 79° 32.344' E) respectively at different elevation gradient, from the altitudinal range of 500m to 2100m above mean sea level in Nainital district.

Each forest study was carried out into three forest site i.e. hill top, hill mid and hill base in open, moderate and

close canopy in the present study to assess the vegetation analysis and standing tree biomass. Each forest represents varied species composition and structure. There exists a varied topography, soil conditions, and forest quality from lower elevation to higher elevation representing Sal forest following subtropical pine forest and Oak forest moving toward the higher elevation in Nainital district of Kumaun Himalayas.




Fig. 1. Description of forest sites in Nainital district of Kumaun Himalaya.

Source: www.mapofindia.com and Google earth.

**Oak forest:** Area falls under moist temperate forest type (Lesser Himalayan Zone: 1800- 2700 m) according to revised forest classification of India, (Champion and Seth, 1968). Oak dominated forest is mainly found associated with *Quercus floribunda*, *Cedrus deodara*, *Cupressus torulosa*, *Rhododendron arborium*, *Myrica esculenta*, *Alnus nepalensis*, *Acer oblongum*, *Lyonia ovalifolia*, *Cornus oblonga*, *Debrigesia longifolia* etc. in the higher regions Nainital district. Oak is well known temperate tree species for its quality fuel and fodder production in Kumaun Himalaya.

**Pine forest:** Pine was found associated with *Cupressus torulosa*, *Acer oblongum*, *Quercus spp.*, *Myrica esculenta*, *Rhododendron arborium* etc. Pine forest was found extensively distributed in the subtropical regions of Nainital district. Pine, family; Pinaceae, is highly economic species which produce good timber, resin, bedding and packing material and several other non timber forest products.

**Sal forest:** Sal is a semi-evergreen large gregarious tree species and (Outer Himalayan Zone: 500-1250 m) was found generally associated with *Mallatos philipinsis*, *Toona ciliata*, *Holoptela intrifolia*, *Terminalia spp.*, *Bauhinia spp.*, *Syzygium cumini*, *Schleichera oleosa*, *Melia azadirac*, *Murraya koenigii*, *Tectona grandis* etc. Sal forest is widely distributed along the foot hills of Kumaun Himalayan regions. Sal, family; Dipterocarpaceae, is an important species to produce valuable hardwood timber and non timber products like leaves and seed oils. The species reveals economic, social and cultural importance in India.

### Vegetation analysis

For the phytosociological study, the quadrat method was used. The number and size of the quadrats were determined by the running mean method (Kershaw, 1973) and species area curve (Misra, 1968). 30 plots of 10 × 10 m at each site were randomly established for determination of species richness and other vegetation parameters. Circumference at breast height (CBH at 1.37 m from the ground) of individual tree and sapling was measured in each quadrat. Individual trees having ≥30 cm girth were considered as trees. Density, frequency, abundance, basal area and Importance value index (IVI) were calculated following Cottam and Curtis (1956).

### Biomass and carbon stock estimation

The estimation of biomass was carried out using the allometric equation for each tree component as developed by Rawat and Singh (1988), Chaturvedi and Singh (1987), Lodhiyal et al. (1992) Rawat (1983); Rana et al. (1989) and Tewari et al. (1985)

$$\ln Y = a + b \ln X$$

Where,

$\ln$  = natural log,  $Y$  = dry weight of component ( $\text{kg tree}^{-1}$ ),  $X$  = CBH (cm),  $a$  = the y intercept and  $b$  = slope of regression.

Carbon stock was determined by using the biomass value of species multiplied by factor ( $C = \text{Biomass} \times 0.475$ ) as given by (Magnussen and Reed, 2004; Singh and Lodhiyal, 2009; Lodhiyal, 2014; Kapkoti, 2016, Bhakuni, 2016, Brown, 2002; Jharia et al., 2014).

## Results

### Vegetation analysis:

#### Chir-pine forest Site-1 (Sanitorium)

##### Trees

The total six species were present i.e. *Pinus roxburghii*, *Quercus leucotrichophora*, *Myrica esculenta*, *Rhododendron arborium*, *Lyonia obifolia* and *Cupressus torulosa* in this forest site. The total tree density was 517 ind.ha<sup>-1</sup>. Of which, *Pinus roxburghii* accounted for maximum 396 ind.ha<sup>-1</sup> followed by *Quercus leucotrichophora* 56 ind.ha<sup>-1</sup> and *Myrica esculenta* 25 ind.ha<sup>-1</sup>. Total basal area of tree was 45.76 m<sup>2</sup>ha<sup>-1</sup>. Of which *Pinus roxburghii* shared 37.04 m<sup>2</sup>ha<sup>-1</sup> followed by *Quercus leucotrichophora* 4.53 m<sup>2</sup>ha<sup>-1</sup>. The IVI of forest was 299.95. Maximum IVI shared *Pinus roxburghii* 210.71 followed by *Quercus leucotrichophora* 37.75 (Table1).

16m<sup>2</sup>ha<sup>-1</sup>. Total basal area of tree was 49.42m<sup>2</sup>ha<sup>-1</sup>. Of which *Quercus leucotrichophora* shared 45.3 m<sup>2</sup>ha<sup>-1</sup> followed by *Acer oblungum* 1.28m<sup>2</sup>ha<sup>-1</sup>. The IVI of forest was 299.93. Maximum IVI shared *Quercus leucotrichophora* 235.11 followed by *Myrica esculenta* 28.56 (Table 2).

#### Sal forest site 3 (Ranibag)

##### Trees

The total six species were present i.e. *Shorea robusta*, *Syzygium cumini*, *Mallotos Phillippensis*, *Toona ciliata*, *Emblia officinailis*, *Tectona grandis* and *Terminelia elliptica* in this forest site. The total tree density was 796ind.ha<sup>-1</sup>. Of which, *Shorea robusta* accounted for maximum 620 ind.ha<sup>-1</sup> followed by *Syzygium cumini* 68 ind.ha<sup>-1</sup> and *Mallotos Phillippensis* 32 ind.ha<sup>-1</sup>. Total basal area of tree was 40.41m<sup>2</sup>ha<sup>-1</sup>. Of which *Shorea robusta* shared 38.8m<sup>2</sup>ha<sup>-1</sup> followed by *Syzygium cumini* 0.72 m<sup>2</sup>ha<sup>-1</sup>. The IVI of forest was 299.8. Maximum IVI shared *Shorea robusta* 216.15 followed by *Syzygium cumini* 30.654 (Table 3).

##### Tree biomass

#### Pine forest (Sanitorium Site-1)

Total biomass of tree species was 135.43 tha<sup>-1</sup>. The maximum biomass account for *Pinus roxburghii* 51.88 tha<sup>-1</sup> followed by

**Table 1.** Vegetation analysis of trees in *Pinus roxburghii* forest in site-1.

Name of species	Density (ind.ha <sup>-1</sup> )	Frequency (%)	Abundance	A\F Ratio	TBA (m <sup>2</sup> ha <sup>-1</sup> )	IVI
<i>Pinus roxburghii</i>	396	100	3.96	0.039	37.04	210.71
<i>Quercus leucotrichophora</i>	56	32	1.75	0.054	4.53	37.75
<i>Myrica esculenta</i>	25	24	1	0.041	1.09	19.97
<i>Rhododendron arborium</i>	20	16	1.25	0.078	1.86	16.43
<i>Lyonia obifolia</i>	12	8	1.5	0.187	0.85	8.42
<i>Cupressus torulosa</i>	8	8	1	0.125	0.39	6.64
<b>Total</b>	<b>517</b>	<b>188</b>	<b>10.46</b>	<b>0.524</b>	<b>45.76</b>	<b>299.95</b>

**Note:** TBA= Total Basal Area, A/F= Abundance/Frequency and IVI=Important Value Index

#### Oak Forest Site -2 (Kailakhan)

##### Trees

The total five species were present i.e. *Quercus leucotrichophora*, *Myrica esculenta*, *Rhododendron arborium*, *Cedrus deodara* and *Acer oblungum* in this forest site. The total tree density was 424 ind.ha<sup>-1</sup>. Of which, *Quercus leucotrichophora* accounted for maximum 356 ind.ha<sup>-1</sup> followed by *Myrica esculenta* 32m<sup>2</sup>ha<sup>-1</sup> and *Rhododendron arboretum*

*Quercus leucotrichophora* 38.8 tha<sup>-1</sup>. Among the various tree components, biomass of bole component shared 53.08 tha<sup>-1</sup> while stump root contributed 53.08 tha<sup>-1</sup> in Pine forest (Table 4).

#### Oak forest (Kailakhan site-2)

Total biomass of tree species was 337.3 tha<sup>-1</sup>. The maximum biomass account for *Quercus leucotrichophora* 310.8 tha<sup>-1</sup> followed by *Myrica esculenta* 11.94 tha<sup>-1</sup>. Among the various

**Table 2.** Vegetation analysis of trees in *Quercus leucotrichophora* forest in site-2.

Name of Species	Density (ind.ha <sup>-1</sup> )	Frequency( %)	Abundance	A\F Ratio	TBA(m <sup>2</sup> ha <sup>-1</sup> )	IVI
<i>Quercus leucotrichophora</i>	356	100	3.56	0.03	45.3	235.11
<i>Myrica esculenta</i>	32	32	1	0.03	0.98	28.56
<i>Rhododendron arborium</i>	16	16	1	0.06	1.08	15.48
<i>Cedrus deodara</i>	8	8	1	0.12	0.78	8.22
<i>Acer oblungum</i>	12	12	1	0.08	1.28	12.56
<b>Total</b>	<b>424</b>	<b>168</b>	<b>7.56</b>	<b>0.32</b>	<b>49.42</b>	<b>299.93</b>

**Note:** TBA= Total Basal Area, A/F= Abundance/Frequency, IVI=Important Value Index

**Table 3.** Vegetation analysis of trees in *Shorea robusta* forest in site-3.

Name of Species	Density (ind.ha <sup>-1</sup> )	Frequency( %)	Abundance	A\F Ratio	TBA(m <sup>2</sup> ha <sup>-1</sup> )	IVI
<i>Shorea robusta</i>	620	100	6.2	0.062	38.80	216.15
<i>Syzygium cumini</i>	68	48	1.41	0.029	0.72	30.65
<i>Mallotus philippensis</i>	32	32	1	0.031	0.21	18.09
<i>Toona ciliata</i>	16	16	1	0.062	0.06	8.93
<i>Emblia officinalis</i>	8	8	1	0.125	0.17	4.82
<i>Tectona grandis</i>	28	16	1.75	0.109	0.37	11.19
<i>Terminelia elliptica</i>	24	16	1.5	0.093	0.08	9.97
<b>Total</b>	<b>796</b>	<b>236</b>	<b>13.86</b>	<b>0.51</b>	<b>40.41</b>	<b>299.80</b>

**Note:** TBA= Total Basal Area, A/F= Abundance/Frequency, IVI=Important Value Index

**Table 4.** Component wise tree biomass (t ha<sup>-1</sup>) of each tree species in forest site 1.

Name of the species	Bole	Branch	Foliage	Twig	Leaf	Stump root	Lateral root	Fine root	Total
<i>Pinus roxburghii</i>	25.13	18.8	0.06	-	-	6.12	1.58	0.19	51.88
<i>Quercus leucotrichophora</i>	16.94	10.28	-	3.87	2.26	4.89	0.53	0.03	38.8
<i>Myrica esculenta</i>	4.74	2.97	-	2.66	0.71	1.06	0.17	0.01	12.32
<i>Rhododendron arboreum</i>	1.65	1.05	-	0.36	0.14	1.02	0.34	0.002	4.56
<i>Lyonia ovalifolia</i>	3.3	20.1	-	0.77	0.45	0.98	0.1	0.007	25.70
<i>Cupressus</i>	1.32	0.05	0.18	0.04	-	0.335	0.2	0.04	2.16
<b>Total</b>	<b>53.08</b>	<b>53.25</b>	<b>0.24</b>	<b>7.7</b>	<b>3.56</b>	<b>14.40</b>	<b>2.92</b>	<b>0.279</b>	<b>135.43</b>

**Table 5.** Component wise tree biomass (t ha<sup>-1</sup>) of each tree species in forest site 2.

Name of the species	Bole	Branch	Twig	Leaf	Stump root	Lateral root	Fine root	Total
<i>Quercus leucotrichophora</i>	157.68	94.72	25.38	8.36	10.59	13.07	1.08	310.88
<i>Myrica esculenta</i>	4.85	3.02	1.34	0.78	1.76	0.18	0.01	11.94
<i>Rhododendron arborium</i>	1.18	0.76	0.27	0.11	0.71	0.23	0.03	3.29
<i>Cedrus deodara</i>	0.48	-	-	-	0.80	-	-	1.28
<i>Acer oblungum</i>	4.43	2.65	0.95	0.56	1.19	0.13	0.008	9.92
<b>Total</b>	<b>168.62</b>	<b>101.15</b>	<b>27.94</b>	<b>9.81</b>	<b>15.05</b>	<b>13.61</b>	<b>1.12</b>	<b>337.32</b>

tree components, biomass of bole component shared 168 tha<sup>-1</sup> while stump root contributed 15.05 tha<sup>-1</sup> in oak forest (Table 5).

### Sal forest (Ranibag site-3)

Total biomass of tree species was 513.20 tha<sup>-1</sup>. The maximum biomass account for *Shorea robusta* 505.08 tha<sup>-1</sup> followed by

**Table 6.** Component wise tree biomass (t ha<sup>-1</sup>) of each tree species in forest site 3.

Name of the Species	Bole	Branch	Twig	Leaf	Root	Total
<i>Shorea robusta</i>	257.83	67.58	30.43	21.23	128	505.08
<i>Syzigium cumini</i>	1.96	0.66	0.30	0.20	0.34	3.46
<i>Mallotus phillippensis</i>	0.41	0.19	0.04	0.02	0.07	0.75
<i>Toona ciliata</i>	0.14	0.06	0.028	0.02	0.03	0.29
<i>Emblia officinailis</i>	0.55	0.16	0.62	0.04	0.07	1.45
<i>Tectona grandis</i>	1.04	0.34	0.139	0.09	0.16	1.79
<i>Terminalia elliptica</i>	0.18	0.07	0.037	0.02	0.04	0.36
<b>Total</b>	<b>262.11</b>	<b>69.08</b>	<b>31.60</b>	<b>21.65</b>	<b>128.74</b>	<b>513.20</b>

**Table 7.** Component wise carbon stock (t C ha<sup>-1</sup>) of each tree species in forest site 1.

Name of the species	Bole	Branch	Foliage	Twig	Leaf	Stump root	Lateral root	Fine root	Total
<i>Pinus roxburghii</i>	11.936	8.93	0.028	-	-	2.907	0.750	0.09	24.64
<i>Quercus leucotrichophora</i>	8.046	4.883	-	1.838	1.073	2.322	0.251	0.01	18.43
<i>Myrica esculenta</i>	2.251	1.410	-	1.263	0.337	0.503	0.080	0.004	5.852
<i>Rhododendron arboreum</i>	0.783	0.498	-	0.171	0.066	0.484	0.161	0.0009	2.16
<i>Lyonia ovalifolia</i>	1.567	9.547	-	0.365	0.213	0.465	0.047	0.0033	12.21
<i>Cupressus</i>	0.627	0.023	0.085	0.019	0	0.159	0.095	0.019	1.028
<b>Total</b>	<b>25.213</b>	<b>25.293</b>	<b>0.114</b>	<b>3.657</b>	<b>1.691</b>	<b>6.842</b>	<b>1.387</b>	<b>0.132</b>	<b>64.33</b>

**Table 8.** Component wise carbon stock (t C ha<sup>-1</sup>) of each tree species in forest site 2.

Name of the species	Bole	Branch	Twig	Leaf	Stump root	Lateral root	Fine root	Total
<i>Quercus leucotrichophora</i>	74.89	44.99	12.05	3.971	5.030	6.208	0.513	147.66
<i>Myrica esculenta</i>	2.303	1.435	0.636	0.370	0.836	0.085	0.004	5.67
<i>Rhododendron arborium</i>	0.560	0.361	0.128	0.052	0.337	0.109	0.014	1.56
<i>Cedrus deodara</i>	0.231	-	-	-	0.380	-	-	0.61
<i>Acer oblongum</i>	2.104	1.259	0.451	0.266	0.565	0.061	0.003	4.71
<b>Total</b>	<b>80.098</b>	<b>48.048</b>	<b>13.271</b>	<b>4.659</b>	<b>7.149</b>	<b>6.464</b>	<b>0.535</b>	<b>160.22</b>

**Table 9.** Component wise carbon stock (t C ha<sup>-1</sup>) of each tree species in forest site 3.

Name of the species	Bole	Branch	Twig	Leaf	Root	Total
<i>Shorea robusta</i>	122.469	32.100	14.45	10.088	60.8	239.91
<i>Syzigium cumini</i>	0.931	0.314	0.142	0.095	0.161	1.64
<i>Mallotus phillippensis</i>	0.194	0.091	0.022	0.011	0.037	0.35
<i>Toona ciliata</i>	0.066	0.030	0.013	0.009	0.017	0.13
<i>Emblia officinailis</i>	0.261	0.080	0.294	0.020	0.034	0.69
<i>Tectona grandis</i>	0.497	0.162	0.066	0.047	0.080	0.85
<i>Terminalia elliptica</i>	0.085	0.033	0.017	0.012	0.023	0.17
<b>Total</b>	<b>124.50</b>	<b>32.814</b>	<b>15.011</b>	<b>10.284</b>	<b>61.153</b>	<b>243.77</b>

*Syzigium cumini* 3.46  $\text{tha}^{-1}$ . Among the various tree components, biomass of bole component shared 262.11  $\text{tha}^{-1}$  while root contributed 128.74  $\text{tha}^{-1}$  in Pine forest (Table 6).

### Carbon stock

#### Pine forest (Sanitorium-site 1)

The total carbon in forest site-1 was 64.33  $\text{t C ha}^{-1}$ . Of this, *Pinus roxburghii* accounted maximum for 24.64  $\text{t C ha}^{-1}$ . Of the total carbon, bole wood component accounted for 25.213  $\text{t ha}^{-1}$  followed by branch in above ground biomass 25.293  $\text{t C ha}^{-1}$  while stump root accounted 6.842  $\text{t C ha}^{-1}$  in belowground part at forest site 1 (Table 7).

#### Oak forest (Kailakhan-site 2)

The total carbon in forest site-2 was 160.22  $\text{t C ha}^{-1}$ . Of this, *Quercus leucotrichophora* accounted maximum for 24.64  $\text{t C ha}^{-1}$ . Of the total carbon, bole wood component accounted for 25.213  $\text{t ha}^{-1}$  followed by branch in aboveground biomass 25.293  $\text{t C ha}^{-1}$  while stump root accounted 6.842  $\text{t C ha}^{-1}$  in belowground part at forest site 1 (Table 8).

#### Sal forest (Ranibag-site 3)

The total carbon in forest site-3 was 243.77  $\text{t C ha}^{-1}$ . Of this, *Shorea robusta* accounted maximum for 239.91  $\text{t C ha}^{-1}$ . Of the total carbon, bole wood component accounted for 124.50  $\text{t ha}^{-1}$  followed by branch in above ground biomass 32.814  $\text{t C ha}^{-1}$  while root accounted 61.153  $\text{t C ha}^{-1}$  in belowground part at forest site 3 (Table 9).

### Discussion

Forest plays a significance role in the development, livelihood and climate they provide timber, non-timber products agricultural implements to hill people and also improve fertility of the agriculture soil and control erosion and forests composition in the hill region, vary from place to place because of altitude, climate, slope, aspect and soil characteristics. Forests of Himalaya play vital role in the sustainable development of forests and mitigation of the climate change. This study provides a glimpse to view insight about, vegetation structure, biomass and carbon stock present in the sites and its contribution in forests of Kumaun Himalayan region. The objectives of study were to assess the vegetation structure

(density, frequency, abundance, basal area and IVI), biomass and carbon content in Pine, Oak and Sal forest.

### Oak forest

The density of oak forest site was 424  $\text{ind.ha}^{-1}$  Of this, present values fall within range 420-1640  $\text{ind.ha}^{-1}$  (Saxena and Singh, 1982) reported for temperate forests of western Himalayan and was higher than 570-760  $\text{ind.ha}^{-1}$  reported for oak forest (Rawat and Singh, 1988) and lower than 550-1250  $\text{ind.ha}^{-1}$  in oak dominated forests in Kumaun Himalaya (Singh, Tamta Tewari and Ram, 2014). The shrub density was observed 668  $\text{ind.ha}^{-1}$ . However, Basal area was 40.41  $\text{m}^2 \text{ha}^{-1}$  which was higher than 33.89-36.83  $\text{m}^2 \text{ha}^{-1}$  reported for (Rawat and Singh, 1988) and lower side than 58.66 – 93.00  $\text{m}^2 \text{ha}^{-1}$  reported for natural forest (Lodhiyal *et al.*, 2014).

The total biomass was reported 337.32  $\text{t ha}^{-1}$  which falls within the range 285-458  $\text{t ha}^{-1}$  reported for oak dominated forest (Rawat and Singh, 1988) and 236-400  $\text{t ha}^{-1}$  for oak dominated forest at higher altitude (Adhakari, Rawat and Singh 1995) but lower than 651-718  $\text{t ha}^{-1}$  of natural forest of Kumaun Himalayans. (Lodhiyal and Lodhiyal, 2014) and (Bhakuni, 2014). The carbon stock was 160.22  $\text{t ha}^{-1}$  is higher than 229-270  $\text{t ha}^{-1}$  reported in oak forest (Lal and Lodhiyal, 2016), 229-341  $\text{t ha}^{-1}$  of natural forest of Kumaun Central Himalaya (Lodhiyal *et al.* 2014).

### Pine forest

Tree density of pine forest was 517  $\text{ind.ha}^{-1}$ . Of this, present values fall within range 420-1640  $\text{ind.ha}^{-1}$  reported for temperate forests of western Himalaya (Saxena and Singh 1982) and was lower than 720  $\text{ind.ha}^{-1}$  reported for *Pinus roxburghii* forest by (Lodhiyal *et al.*, 2013) and lower than 540-1630  $\text{ind.ha}^{-1}$  for pine forest in Kumaun region (Chaturvedi and Singh, 1987). Shrub density was 486  $\text{ind.ha}^{-1}$ . Total basal area 45.77  $\text{m}^2 \text{ha}^{-1}$  was lower than 58.66-93.00  $\text{m}^2 \text{ha}^{-1}$  reported for natural forests (Lodhiyal *et al.*, 2014) on the other side higher side than 25-47  $\text{m}^2 \text{ha}^{-1}$  reported for pine forests in Kumaun region (Chaturvedi and Singh, 1987).

The total biomass estimates was 135  $\text{t ha}^{-1}$  which falls lower than 154-301  $\text{tha}^{-1}$  in pine forests of Kumaun Himalaya (Lal and Lodhiyal, 2015). However Present values fall within the

range 112-283 t ha<sup>-1</sup> reported for pine forests of central Himalaya (Chaturvedi and Singh, 1987). The carbon stock of the study site was 64.33 t ha<sup>-1</sup> on lower side than 341-228 t ha<sup>-1</sup> reported for natural forests (Lodhiyal *et al.*, 2014), and it falls on higher than 97-207 t ha<sup>-1</sup> reported for pine and mix Banj-oak forests (Rana *et al.*, 1989) and 122-86 t ha<sup>-1</sup> for chir-pine van Panchayat forest (Jeena *et al.*, 2008). But present values are on higher side than 59.41 t ha<sup>-1</sup> oak and pine mixed forest of Lohaghat in Kumaun Himalaya (Lodhiyal and Lodhiyal, 2012) 16.73-18.54 t ha<sup>-1</sup> of oak and pine forests in degraded forests (Jeena, Sah, Bhatt and Rawat, 2008).

### Sal forest

Tree density of the forest site was 796 ind.ha<sup>-1</sup> the value of density is greater than the value obtained by (Bandhu, 1969) reported density 200 ind.ha<sup>-1</sup> of northon tropical dry deciduous forest near Baranasi. The plant density in the present study is also higher than the value 183 ind.ha<sup>-1</sup> by (Kumar *et al.*, in 1994) in dry peninsular sal forest similarly value was also higher than 194 ind.ha<sup>-1</sup> reported for sal forest by (Rana *et al.*, 1988). Shrub density was 2056 ind.ha<sup>-1</sup> Total basal area was 40.41 m<sup>2</sup> ha<sup>-1</sup> which is on higher side than 26.7 m<sup>2</sup> ha<sup>-1</sup> as reported by (Rana *et al.*, 1988). However the value of basal area falls lower than 76 m<sup>2</sup> ha<sup>-1</sup> as estimated in the sal forest of Royal Chitwan National Park, Nepal (Sejuwal, 1994).

The total biomass estimates was 513.20 t ha<sup>-1</sup> which falls lower than 330.87 t ha<sup>-1</sup> by (Giril, Aryal, Bhattaraiz, Ghimire, Shrestha and Jha, 1999) sal forests in the Royal Bardia National Park, Nepal. The Carbon stock of tree was 243.77 t ha<sup>-1</sup>, the value of the carbon falls in the range of 123.15 t ha<sup>-1</sup> and 384.20 t ha<sup>-1</sup> by (Pandey and M. Bhusal) in Sal (*Shorea robusta*) forest in two different ecological regions of Nepal.

### Acknowledgement

Authors are thankful to the department of Forestry and Environmental Science, D.S.B. Campus, Kumaun University, Nainital. We also thank Prof. L.S. Lodhiyal, Department of Forestry and Environmental Science, D.S.B. Campus, Kumaun University, Nainital, for reviewing the manuscript and providing

valuable suggestions. We are thankful to Head department of Forestry and Environmental Science, D.S.B. Campus, Kumaun University, Nainital for giving valuable suggestions.

### References

- Adhikari BS, Rawat YS and Singh SP. 1995.** Structure and Function of Altitude Forests of Central Himalaya I. Dry Matter Dynamics. *Annals of Botany*. 75: 237-248.
- Bandhu D. 1969.** A study of the productive structure of northern tropical dry deciduous forest near Vsransi. I. Stand structure and non –photosynthetic biomass. *Tropical Ecology*. 11: 80-102.
- Brown S. 2002.** Measuring, monitoring and verification of carbon benefits for forest-based projects. *Phil. Trans. R. Soc. Lond.* A360: 1669-1684.
- Champion HG and Seth SK. 1968.** *The forest types of India*. Govt. of India, New Delhi.
- Chaturvedi OP and Singh JS. 1987.** The structure and function of pine forest in Central Himalaya: Dry matter dynamics. *Annals of Botany*. 60: 237-252.
- Cottam G and Curtis JT. 1956.** The use of distance measures in phytosociological sampling. *Ecol.* 35: 451-460.
- Curtis JT and McIntosh RP. 1950.** The interrelations of certain analytic and synthetic phytosociological characters. *Ecol.* 31: 434-455.
- Eckholm EP. 1975.** The deterioration of mountain environments. *Science*. 198: 764-770.
- Singh JS and Singh SP. 1992.** Forest of Himalaya: Structure, functioning and impact of man. Gynodaya Prakashan, Nainital, India.
- Jeena BS, Sah P, Bhatt MD and Rawat YS. 2008.** Estimation carbon sequestration rate and total carbon stock in degraded and non degraded sites of Oak and Pine forest of Kumaun central Himalaya. *Ecological society (ECOS), Nepal Ecoprint*. 15: 75-81.
- Jhariya MK, Bargali SS, Swamy SL, Kittur B, Bargali K and Pawar GV. 2014.** Impact of forest fire on biomass and Carbon storage pattern of Tropical Deciduous Forests in Bhoramdeo Wildlife Sanctuary.



- Kershaw KA. 1973.** Quantitative and dynamic plant ecology. Edward Arnold Ltd. London. Pp. 308.
- Kumar R, Singh AK and Abbas G. 1994.** Change in population structure of some dominant trees species of dry peninsular Sal forest. *Indian Forester*. 120: 549-557.
- Lal B and Lodhiyal LS. 2015.** Vegetation structure, biomass and carbon content in *Pinus roxburghii* Sarg. Dominant forests of Kumaun Himalaya. *Environment & We An International Journal of Science & Technology*. 10: 117-124.
- Lal B and Lodhiyal LS. 2016.** Stand Structure, Productivity and Carbon Sequestration Potential of Oak Dominated Forests in Kumaun Himalaya. *Current World Environment*. 11(2): 466-476.
- Lodhiyal LS, Lodhiyal N and Kapkoti B. 2013.** Structure and diversity of tree species in natural forest of Kumaun Himalaya in Uttarakhand *Journal of plant development science*. 5(2): 97- 105.
- Lodhiyal LS, Lodhiyal N and Bhakuni N. 2014.** Standing tree biomass and carbon content in natural forests of Kumaun in central Himalaya. *Journal of plant development Science*. 6(2): 167-174.
- Lodhiyal LS. 2011.** Biodiversity Conservation And Management of Forests: The significance of participation of stakeholders and communities, *Quest- The Journal of UGC-ASC Nainital*. 5(1): 49-81.
- Lodhiyal N and Lodhiyal LS. 2012.** Tree layer composition and carbon content of Oak and Pine in Lohaghat forest of Kumaun Himalaya. *Journal of plant development Science*. 4: 55-62.
- Misra R. 1968.** Ecology Work Book Oxford Publishing Company, Calcutta.
- Phillips EA. 1959.** Methods of vegetation study. Henry Hill & Co. Inc., U.S.A.
- Ram J, Kumar A and Bhatt J. 2004.** Plant diversity in six forest types of Uttaranchal, Central Himalaya, India. *Current Science*. 86: 975-978.
- Rana BS, Singh RP and Singh SP. 1989.** Carbon and energy dynamic of seven Central Himalayan forests. *Tropical Ecology*. s30 (2): 253-264.
- Rana BS, Singh SP and Singh RP. 1989.** Biomass and net primary Productivity in Central Himalaya forests along an altitudinal gradient. *Forest Ecology and Management*. 27: 199-218.
- Rana BS, Singh SP and Singh RP. 1988.** Biomass and productivity of central Himalayan Sal forest. *Tropical Ecology*. 29: 1-6.
- Rawat YS and Singh JS. 1988.** Structure and function of Oak forests in Central Himalaya. Dry matter dynamics. *Annals of Botany*. 62: 397-411.
- Sati VP. 2005.** Natural Resource Conditions and Economic Development in the Uttranchal Himalaya, India. *J. Mt. Sci*. 2(4): 336-350.
- Saxena AK and Singh JS. 1982.** A Phytosociological analysis of woody species in forest communities of a part of Kumaun Himalaya. *Vegetation*. 50(1): 3-22.
- Sejuwal M. 1994.** Above ground biomass estimation in tropical forest of Royal Chitwan National Park (RCNP) M.Sc. Thesis, Central Department of Botany, Tribhuwan University Kathmandu Nepal.
- Singh JS. 2006.** Sustainable development of the Indian Himalayan region: Linking ecological and economic concerns. *Curr. Sci*. 90(6): 784-788.
- Singh JS and Singh SP. 1992.** Forest of Himalaya, Structure, Functioning and impact of Man. Gyanodaya Prakashan, Nainital, India.
- Singh N, Tamata K, Tewari A and Ram J. 2014.** Studies on Vegetational analysis and regeneration status of *Pinus roxburghii* Roxb. and *Quercus leucotrichophora* forest of Nainital forest division. *Global Journal of Science Frontier Research: C Biological Science*. 14(3): 40-47.
- Swank WT and Schreuder HT. 1974.** Comparison of three methods of estimating surface area and biomass for a forest of young eastern white pine. *Forest Science*. 20: 91-100.
- Valdiya KS. 1980.** Stratigraphic scheme of the sedimentary units of the Kumaun Lesser Himalaya. In: K.S. Valdiya and S.B. Bhatia (eds.). *Stratigraphy and Correlation of the Lesser Himalaya Formations* Hindustan Publications Corporation, New York, USA.

**Valdiya KS. 1980.** Geology of Kumaun Lesser Himalaya. Wadia Institute of Himalayan Geology, Dehradun.

**Pandey HP and Bhusal M. 2016.** A comparative study on carbon stock in Sal (*Shorea robusta*) forest in two different ecological regions of Nepal. *Banko Janakari*. 26(1): 24-31.

**Giri B, Aryal B. Bhattarai, Ghimire SK, Shrestha KK and Jha PK. 1999.** Vegetation Composition, Biomass Production and Regeneration in *Shorea robusta*: Forests in the Royal bardia national park, Nepal. *Nepal Journal of Science and Technology*. (1) 47-56.